

MATH 1300 ASSIGNMENT PROBLEMS (UNIT 4)

[10] 1. Let $A = \begin{bmatrix} 5 & 2 & 4 \\ 2 & -3 & 5 \\ -3 & 2 & 1 \end{bmatrix}$ and let $B = \begin{bmatrix} 3 & 2 & -4 \\ 5 & -1 & 3 \\ 2 & 4 & 0 \end{bmatrix}$. Find the following.

(a) $2A + B^T$.

(b) AB

(c) BA

(d) The matrix C for which $A - C^T = 2B$.

[10] 2.(a) Which of the following matrices are elementary matrices?

(i) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 3 & 0 & 1 \end{bmatrix}$

(ii) $\begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{bmatrix}$

(iii) $\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$

(b) Let $A = \begin{bmatrix} 1 & 3 & 2 \\ 2 & 5 & 4 \\ 3 & 1 & 2 \end{bmatrix}$. Find an elementary matrix E such that $EA = B$ if

(i) $B = \begin{bmatrix} 1 & 3 & 2 \\ 2 & 5 & 4 \\ 5 & 7 & 6 \end{bmatrix}$

(ii) $B = \begin{bmatrix} 2 & 5 & 4 \\ 1 & 3 & 2 \\ 3 & 1 & 2 \end{bmatrix}$

(iii) $B = \begin{bmatrix} 1 & 3 & 2 \\ 2 & 5 & 4 \\ 6 & 2 & 4 \end{bmatrix}$

(c) Let $A = \begin{bmatrix} 3 & 1 \\ 1 & 0 \end{bmatrix}$. Find two elementary matrices E_1 and E_2 such that $E_2 E_1 A = I$.

[10] 3. Find the inverse of $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 2 & 3 & 3 & 3 \\ 3 & 4 & 5 & 5 \\ 4 & 5 & 6 & 7 \end{bmatrix}$. Show all your work and verify that your answer

is correct.

[10] 4. Consider the following system of linear equations

$$\begin{aligned}x_1 + x_2 + x_3 + x_4 &= 3 \\2x_1 + 3x_2 + 3x_3 + 3x_4 &= 4 \\3x_1 + 4x_2 + 5x_3 + 5x_4 &= 1 \\4x_1 + 5x_2 + 6x_3 + 7x_4 &= 2\end{aligned}$$

(a) Rewrite this system of linear equations as a single matrix equation in the form $\mathbf{Ax} = \mathbf{b}$.

(b) Use the matrix \mathbf{A}^{-1} to find the solution \mathbf{x} . [Hint: See problem 3 for \mathbf{A}^{-1} .]

[10] 5. Provide examples to illustrate the following.

(a) A 3×3 non zero matrix \mathbf{B} such that $\mathbf{B}^2 = \mathbf{O}$.

(b) A 3×3 matrix \mathbf{C} such that $\mathbf{C}^2 = \mathbf{C}$ with $\mathbf{C} \neq \mathbf{I}$ and $\mathbf{C} \neq \mathbf{O}$.

(c) A 3×3 matrix \mathbf{D} such that $\mathbf{D}^T = \mathbf{D}$ with $\mathbf{D} \neq \mathbf{I}$ and $\mathbf{D} \neq \mathbf{O}$.

(d) A 3×3 matrix \mathbf{F} such that $\mathbf{F}^T = -\mathbf{F}$ with $\mathbf{F} \neq \mathbf{O}$.

[10] 6. The citizens of Oz have a choice of 3 political parties in their municipal elections, the Blue party, the Green party or the Red party. A study of past voting patterns shows that if a citizen voted for the Blue party in one election, the probability that he/she will vote for the Blue party in the next election is 80%, the probability he/she will vote for the Green party is 10% and the probability he/she will vote for the Red party is 10%. If a citizen voted for the Green party in one election, the probability that he/she will vote for Green party in the next election is 70%, the probability he/she will vote for the Blue party is 20% and the probability that he/she will vote for the Red party is 10%. If a citizen voted for the Red party in one election, the probability that he/she will vote for the Red party in the next election is 60%, the probability that he/she will vote for the Blue party is 20% and the probability that he/she will vote for the Green party is 20%.

(a) Find the transition matrix for the voting intentions of the citizens of Oz.

(b) If the vote distribution at the last election was Blue 40%, Green 35%, Red 25%, find the probable vote distribution at the next election.

(c) Find the long term steady state distribution of the votes in Oz.